

“Weakly” Elliptic Gorenstein Singularities of Surfaces

András Némethi¹

Department of Mathematics, The Ohio State University, Columbus OH 43210, USA,
(e-mail: nemethi@math.ohio-state.edu)

1. Introduction.

Let p be a singularity of a normal two-dimensional analytic surface X . In general $\pi : M \rightarrow X$ will denote a resolution of (X, p) , and $A = \pi^{-1}(p)$. It is well-known that the dual resolution graph of the germ (X, p) determines the topology of the germ (X, p) . Indeed, the (real) link of (X, p) can be reconstructed from the resolution graph as a plumbing. Conversely, by a result of W. Neumann [15], the topology of (X, p) determines the (minimal) resolution graph. Therefore, if an invariant of (X, p) depends only on the dual resolution graph, we say that it is a “topological invariant”. For example, the self-intersection Z_{num}^2 of Artin’s (fundamental) cycle Z_{num} , or the Euler-characteristic $\chi(D)$ of any cycle D supported by A , can be determined from the graph.

Now, it is fascinating to investigate if an invariant, a priori defined from the analytic structure of (X, p) , is topological or not. In this article, we ask this question for the geometric genus $p_g = h^1(M, \mathcal{O}_M)$ and the Hilbert–Samuel function of (X, p) , in particular for the multiplicity $mult(X, p)$ and the embedding dimension $embdim(X, p) = \dim m_p/m_p^2$. In general, these invariants are not topological, but if we restrict our study to some special classes, then they can be determined from the graph.

The first result of this type was obtained by M. Artin [1, 2] for rational singularities. He proved that they can be characterized topologically:

$$(X, p) \text{ is rational, i.e. } p_g = 0 \Leftrightarrow \chi(Z_{num}) = 1 \Leftrightarrow \min_{D>0} \chi(D) \geq 1.$$

Moreover, for these singularities, the Hilbert–Samuel function (hence $mult(X, p)$ and $embdim(X, p)$ too) can also be computed from the resolution graph.

The elliptic singularities were introduced by P. Wagreich in [21] (cf. “Terminology” at the end of the introduction). They are defined topologically:

$$(X, p) \text{ is elliptic} \Leftrightarrow \chi(Z_{num}) = 0 \Leftrightarrow \min_{D>0} \chi(D) = 0.$$

The class of elliptic singularities contains all the singularities with $p_g = 1$, and all the Gorenstein singularities with $p_g = 2$; but an elliptic singularity can have arbitrary high geometric genus (see e.g. the examples after Theorem C). The next step in the above program was obtained by H. Laufer [12]. He proved that the Gorenstein singularities with $p_g = 1$ (he called them minimally elliptic singularities) can be characterized topologically (cf. 2.7), and for these singularities all the above (a priori) analytical invariants are topological.

¹Partially supported by NSF Grant No. DMS-9622724

Moreover, he noticed that singularities with $p_g = 1$ (without Gorenstein assumption), or even Gorenstein singularities with $p_g = 2$ (or $p_g \geq 2$) cannot be characterized topologically. In this second case, one can easily construct pairs of hypersurface singularities with the same resolution graph but different p_g (cf. 2.22). But in all these examples (known by the author) either there is a non-rational exceptional divisor in the resolution, or the graph is not a tree, i.e. $H^1(A, \mathbf{Z}) \neq 0$. (Notice that $H^1(A, \mathbf{Z}) = 0$ if and only if the link of (X, p) is a rational homology sphere.)

The main message of the present paper is that for Gorenstein singularities with $H^1(A, \mathbf{Z}) = 0$ the Artin–Laufer program can be continued. Here we give the complete answer in the case of elliptic singularities.

Elliptic singularities were intensively studied by Wagreich [21], Laufer [12], S. S.-T. Yau [23, 24, 25, 26, 27, 28, 29] and others. In Yau’s papers, the fundamental topological invariant is the “elliptic sequence”. In the numerical Gorenstein case its definition is the following. Let $\pi : M \rightarrow X$ be the *minimal* resolution of (X, p) , and $A = \pi^{-1}(p)$ as above. Let Z_K be the canonical cycle (cf. §2). The elliptic sequence consists of a sequence $\{Z_{B_j}\}_{j=0}^m$, where Z_{B_j} is Artin’s (fundamental) cycle of $B_j \subset A$. We define $\{B_j\}_j$ inductively as follows. For $j = 0$ take $B_0 = A$ hence $Z_{B_0} = Z_{\text{num}}$. Then $Z_K \geq Z_{B_0}$. If $Z_K > Z_{B_0}$ then we set $B_1 := |Z_K - Z_{B_0}|$. Similarly, if B_i is already defined for any $i \leq j$, then $Z_K \geq Z_{B_0} + \cdots + Z_{B_j}$ (for details, see 2.10). If the inequality is strict then we define $B_{j+1} := |Z_K - Z_{B_0} - \cdots - Z_{B_j}|$, otherwise we stop. The length of the elliptic sequence $\{Z_{B_j}\}_{j=0}^m$ is $m + 1$. The case $m = 0$ corresponds exactly to the minimally elliptic singularities of Laufer [12].

S. S.-T. Yau proved that for a numerical Gorenstein elliptic singularity $p_g \leq m + 1$ ([29, (3.9)], cf. also 2.19). Particular examples show that strict inequality can occur.

In the first sections we give several characterizations of the “extremal property” $p_g = m + 1$. The most important characterization, from the point of view of the present manuscript, is the following:

Theorem A. *Let (X, p) be an elliptic numerical Gorenstein singularity and set $C'_j = \sum_{i \geq j} Z_{B_i}$. Then $p_g = m + 1$ if and only if the “obstruction line bundles” $\mathcal{O}_{C'_j}(-Z_{B_{j-1}})$ are trivial for all $1 \leq j \leq m$.*

Now, the main technical result of the paper (see section 4) says:

Theorem B. *Let (X, p) be an elliptic Gorenstein singularity. Fix an integer $1 \leq k \leq m$ and assume that for $k + 1 \leq j \leq m$ the “obstruction line bundles” $\mathcal{O}_{C'_j}(-Z_{B_{j-1}})$ are trivial. (If $k = m$ then this assumption is vacuous.) Then $\mathcal{O}_{C'_k}(-Z_{B_{k-1}})$ has finite order. (Actually, its order is not greater than $k!$.)*

Now, if $H^1(A, \mathbf{Z}) = 0$ then $\text{Pic}(D)$ is torsion free for any positive cycle D . Hence, we obtain the main result of the paper:

Theorem C. *Assume that (X, p) is an elliptic Gorenstein singularity with $H^1(A, \mathbf{Z}) = 0$. Then p_g is a topological invariant. In fact $p_g = m + 1 =$ the length of the elliptic sequence in the minimal resolution of (X, p) .*

Actually, one of the starting points of our investigation was S. S.-T. Yau's result in [27], which says that *hypersurface* singularities with $H^1(A, \mathbf{Z}) = 0$ and $p_g = 2$ satisfies $m = 1$ (cf. Remark 4.14). (In Yau's terminology, a numerical Gorenstein elliptic singularity with $p_g = m + 1$ is called "maximally elliptic".)

Examples. [21, 29] Set $(X_i, 0) \subset (\mathbf{C}^3, 0)$ given by $(X_1, 0) = \{z^2 + y^3 + x^{9+6m} = 0\}$, $(X_2, 0) = \{z^2 + y^3 + x^{11+6m} = 0\}$, $(X_3, 0) = \{z^3 + y^3 + x^{4+3m} = 0\}$, $(X_4, 0) = \{z^3 + y^3 + x^{5+3m} = 0\}$ (where $m \geq 0$). Then, in all these cases, $H^1(A_i, \mathbf{Z}) = 0$, $(X_i, 0)$ is elliptic, and $p_g = m + 1 =$ the length of the elliptic sequence ($1 \leq i \leq 4$) (cf. also with 6.4.c).

In sections 5 and 6, we generalize Laufer's results about the minimally elliptic singularities [12] (which corresponds to $m = 0$). Namely, we prove:

Theorem D. *Assume that (X, p) is an elliptic Gorenstein singularity with $p_g = m + 1$. (If $m = 0$ or $H^1(A, \mathbf{Z}) = 0$ then the last assumption is satisfied.) Let Z_{num} be Artin's cycle in the minimal resolution $(M, A) \rightarrow (X, p)$. Then:*

- (a) *If $Z_{num}^2 \leq -2$, then $m_p \mathcal{O}_M = \mathcal{O}(-Z_{num})$, hence $\text{mult}(X, p) = -Z_{num}^2$;*
- (b) *If $Z_{num}^2 = -1$, then $m_p \mathcal{O}_M = m_Q \mathcal{O}(-Z_{num})$ for some smooth point Q of A , and $\text{mult}(X, p) = 2$;*
- (c) *$\text{emb dim}(X, p) = \max(3, -Z_{num}^2)$;*
- (d) *$H^0(M, \mathcal{O}(-kZ_{num})) = m_p^k$ for any $k \geq 0$ provided that $Z_{num}^2 \leq -3$;*
- (e) *$\dim \mathcal{O}_{(X,p)}/m_p^k = \chi(\mathcal{O}_{kZ_{num}}) + 1$, and $\dim m_p^k/m_p^{k+1} = -kZ_{num}^2$ for any $k \geq 1$ provided that $Z_{num}^2 \leq -3$; (for the cases $Z_{num}^2 = -1$ or -2 , see (6.4)).*

Some partial results in this direction were obtained by S. S.-T. Yau in his series of papers (we will cite them in the body of the paper at the corresponding places). In order to have a self-contained presentation, we reprove those facts what we use from his work. In some cases we follow Yau's original arguments, in other cases we give different proofs (original ones or arguments in the spirit of [17]). The article of M. Reid [17] was extremely helpful for the author. Actually, the proofs in sections 5 and 6 have their origins in [17]. One of the ideas of the proof of Theorem B was borrowed from J. Wahl's paper [22]. On the other hand, we emphasize that almost all the classical arguments, used in the case of rational or minimally elliptic singularities, and based on some vanishing theorems or on the numerically 1- or 2-connectivity of Z_{num} , in our general situation fail, and we had to replace them by different arguments.

Some of the results of the present article (especially, the multiplicity and embedding dimension computations) can be compared with the results of U. Karras and J. Stevens proved for Kodaira, respectively Kulikov singularities. For details, see [6, 7] and [19, 20].

Finally, we notice that Theorem D is not true if $\chi(Z_{num}) < 0$. E.g. for $(X, 0) = \{x^3 + y^4 + z^7 = 0\}$ one can verify that $\chi(Z_{num}) = -1$, $H^1(A, \mathbf{Z}) = 0$, and $Z_{num}^2 = -2$, but $\text{mult}(X, 0) = 3$.

Terminology: Singularities characterized by $\chi(Z_{num}) = 0$ sometimes are called "weakly" elliptic. By this terminology one wants to emphasize the difference between these singularities and the "strongly" elliptic singularities, defined by $p_g = 1$; cf. also Yau's papers.

Notice also that in the terminology of M. Reid [17], “elliptic” means “minimally elliptic” in the sense of Laufer (cf. 2.7). In this article we will adopt the terminology used by Wagreich and Laufer.

The author wishes to thank École Polytechnique at Palaiseau, University of Nice and University of Nantes (especially Professors C. Sabbah, M. Merle and F. Elzein) for their hospitality and excellent working atmosphere.

2. Preliminaries.

We fix a normal surface singularity (X, p) . Let $\pi : M \rightarrow X$ be its *minimal* resolution, and let $A = \cup_i A_i$ be the decomposition of the exceptional set A into irreducible components. Let K be the canonical divisor on M , hence:

$$(2.1) \quad A_i K = -A_i^2 - 2 + 2g_i + 2\delta_i \quad \text{for all } i,$$

where g_i is the genus of A_i , and δ_i is the sum of all delta-invariants of the singular points (the “number of nodes and cusps”) on A_i [18].

In this paper all the cycles will be integer combinations of the A_i ’s. If $D = \sum_i n_i A_i$, we write $n_i = m_{A_i}(D)$. $|D|$ denotes the support of D .

We denote Artin’s numerical (fundamental) cycle by Z_{num} , i.e. Z_{num} is the minimal positive cycle Z with $ZA_i \leq 0$ for all i [1, 2]. (X, p) is called numerical Gorenstein if there is a cycle Z_K (called canonical cycle) which satisfies $Z_K A_i = -KA_i$ for all i . If Z_K exists, then (by 2.1) Z_K is trivial if and only if (X, p) is a Du Val singularity. Otherwise $m_{A_i}(Z_K) > 0$ for all A_i (cf. [14], page 490). Moreover, if $Z_K \neq 0$, then from (2.1) one has $Z_K A_i \leq 0$ for all i , hence $Z_{num} \leq Z_K$.

By a (Kodaira type) vanishing theorem (cf. [17], Exercise 15, page 119), for any cycle $D \geq 0$ which satisfies $A_i D \leq 0$ for all A_i , and any line bundle $\mathcal{L}$ with $\deg_{A_i} \mathcal{L} \geq KA_i$ for all i :

$$(2.2) \quad \begin{cases} h^1(M, \mathcal{O}(-D) \otimes \mathcal{L}) = 0, \text{ hence in the numerical Gorenstein case:} \\ H^0(M, \mathcal{O}(-D)) \rightarrow H^0(\mathcal{O}_{Z_K}(-D)) \text{ is onto.} \end{cases}$$

For any cycle D , we denote the Euler–characteristic $h^0(\mathcal{O}_D) - h^1(\mathcal{O}_D)$ by $\chi(D)$. By Riemann–Roch theorem $\chi(D) = -D(D + K)/2$. By [12], (2.6):

$$(2.3) \quad H^0(\mathcal{O}_{Z_{num}}) = \mathbf{C} = \{\text{constants}\}, \text{ hence } \chi(Z_{num}) \leq 1.$$

By a result of Artin [1, 2], the following facts are equivalent:

$$(2.4) \quad (X, p) \text{ is rational (i.e. } p_g = 0) \Leftrightarrow \chi(Z_{num}) = 1 \Leftrightarrow \min_{D \geq 0} \chi(D) \geq 1.$$

The dual resolution graph of a rational singularity is a tree, and all the vertices corresponds to smooth rational curves.

In [21], Wagreich introduced the elliptic singularities. They are defined by the property $\min_{D>0} \chi(D) = 0$. In particular, by (2.3) and (2.4), $\chi(Z_{num}) = 0$. The inverse implication is also true, see e.g. [12] (4.2), hence:

$$(2.5) \quad (X, p) \text{ is elliptic} \Leftrightarrow \chi(Z_{num}) = 0 \Leftrightarrow \min_{D>0} \chi(D) = 0.$$

Elliptic singularities include all the singularities with $p_g = 1$ and all the Gorenstein singularities with $p_g = 2$ (cf. the proof of 4.13); but elliptic singularities can have arbitrary high geometric genus.

Following Laufer (Definition 3.1 [12]), we say that a cycle $E > 0$ is minimally elliptic if $\chi(E) = 0$ and $\chi(D) > 0$ for all cycles $0 < D < E$. Laufer in [12] (3.2) proved that if $\chi(Z_{num}) = 0$ then there exists a unique minimally elliptic cycle E . Hence, in the elliptic numerical Gorenstein case:

$$(2.6) \quad E \leq Z_{num} \leq Z_K.$$

The minimally elliptic singularities are characterized by the following equivalent properties (cf. Laufer's paper [12]) (notice that (b), (c) and (d) are topological properties; (d) provides the name of the singularity):

$$(2.7) \quad \begin{cases} (a) (X, p) \text{ is Gorenstein and } p_g = 1; \\ (b) Z_{num} \text{ is a minimally elliptic cycle;} \\ (c) (X, p) \text{ is numerical Gorenstein with } Z_K = Z_{num}; \\ (d) \chi(Z_{num}) = 0 \text{ and any connected proper subvariety of } A \\ \quad \text{supports a rational singularity.} \end{cases}$$

2.8. If (X, p) is elliptic, then the topology of the irreducible exceptional divisors A_i cannot be very complicated. Wagreich in [21], page 428, proved that precisely one of the followings hold: (a) Precisely one component A_{i_0} satisfies $\chi(A_{i_0}) = 0$ (which is either a smooth elliptic curve or a rational curve with $\delta_{i_0} = 1$) and the other irreducible exceptional divisors are smooth rational curves. (In this case $E = A_{i_0}$). Or (b): all the exceptional divisors are smooth rational curves.

Actually, from the uniqueness of the minimally elliptic cycle it follows that all the connected components of $A \setminus |E|$ support the exceptional set of rational singularities, hence the dual graph of (X, p) can be obtained from the dual graph of $|E|$ by gluing trees whose vertices correspond to smooth rational curves. In particular, the restriction map

$$(2.9) \quad H^1(A, \mathbf{Z}) \rightarrow H^1(|E|, \mathbf{Z}) \text{ is an isomorphism.}$$

In a series of papers S. S.-T. Yau investigated the properties of elliptic singularities (he called them “weakly elliptic”), cf. [23, 24, 25, 26, 27, 28, 29]. His main tools are the “computation sequence” introduced by Laufer in [10] and [12], and the “elliptic sequence”. The interested reader can find in Yau's papers the definition of the elliptic sequence for an arbitrary elliptic singularity. In the non-numerical Gorenstein case the sequence has a lot of anomalies (see e.g. some examples on the page 881 of [28]), but in the numerical

Gorenstein case it is a powerful tool. Since the definition in this later case is easier and more natural, and it is sufficient for our goals, we adopt the general definition for this situation.

2.10. The construction of the elliptic sequence. Consider the minimal resolution of an elliptic numerical Gorenstein singularity. The elliptic sequence consists of the sequence $\{Z_{B_j}\}_{j=0}^m$, where Z_{B_j} is the numerical (fundamental) cycle of $B_j \subset A$. We define $\{B_j\}_j$ inductively as follows. For $j = 0$ take $B_0 = A$ hence $Z_{B_0} = Z_{num}$. By (2.6) $E \leq Z_{num} \leq Z_K$.

If $Z_{B_0} < Z_K$, we define $B_1 := |Z_K - Z_{num}|$. Then B_1 is connected. Indeed, assume that B_1 has more connected components $\{B_{1,t}\}_t$; then write $Z_K - Z_{num} = \sum_t D_t$ with $|D_t| = B_{1,t}$. Then the vanishing $\chi(Z_K - Z_{num}) = 0$ and (2.5) imply that $\chi(D_t) = 0$ for all t . Then by [12] (3.2), each $B_{1,t}$ supports a minimally elliptic cycle. This contradicts the uniqueness of the minimally elliptic cycle E . Now, using again $\chi(Z_{num}) = 0$, one gets $Z_{num}(Z_K - Z_{num}) = 0$. Since $Z_{num}A_i \leq 0$ for all A_i , it follows that for any $A_i \subset B_1$ the equality $Z_{num}A_i = 0$ holds. In particular, by the non-degeneracy of the intersection form, $B_1 \neq B_0$. Moreover, for any $A_i \subset B_1$ one has: $A_i(Z_K - Z_{B_0}) = A_iZ_K$, hence B_1 supports a numerical Gorenstein singularity with canonical cycle $Z_K - Z_{B_0}$. This singularity is again elliptic. Indeed, $\min_{D \subset B_1} \chi(D) \geq \min_{D \subset B_0} \chi(D) = 0$, but $\chi(Z_K - Z_{num}) = 0$, hence $\min_{D \subset B_1} \chi(D) = 0$. In particular, (cf. 2.6), $E \leq Z_{B_1} \leq Z_K - Z_{B_0}$.

Now we repeat the above arguments. If $Z_{B_1} < Z_K - Z_{B_0}$, we define $B_2 := |Z_K - Z_{B_0} - Z_{B_1}|$ and we verify that it supports an elliptic singularity with canonical cycle $Z_K - Z_{B_0} - Z_{B_1}$ and for any $A_i \subset B_2$ the vanishing $A_iZ_{B_1} = 0$ holds. If we repeat this procedure, after a finite step we will obtain $Z_{B_m} = Z_K - Z_{B_0} - \dots - Z_{B_{m-1}}$, i.e. the numerical cycle and the canonical cycle of B_m coincides, hence by (2.7) B_m supports a minimally elliptic singularity with $E = Z_{B_m}$. In particular $B_m = |E|$.

If we contract the connected exceptional curve $B_j \subset M$, we obtain a unique singular point; this will be denoted by $(M/B_j, p_j)$ ($0 \leq j \leq m$). It is convenient to introduce the notations $C_t = \sum_{i=0}^t Z_{B_i}$ and $C'_t = \sum_{i=t}^m Z_{B_i}$ ($0 \leq t \leq m$) too.

2.11. Definition/first properties of the elliptic sequence:

(a) $B_0 = A$, $B_1 = |Z_K - Z_{B_0}|$, $B_2 = |Z_K - Z_{B_0} - Z_{B_1}|$, $\dots$, $B_m = |E|$; each B_j is connected and the inclusions $B_{j+1} \subset B_j$ are strict. Moreover, $Z_{num} = Z_{B_0} \supset Z_{B_1} \supset \dots \supset Z_{B_m} = E$.

(b) If $A_i \subset B_{j+1}$ then $A_iZ_{B_j} = 0$ for all i and j . In particular, $Z_{B_i}Z_{B_j} = 0$ for all $0 \leq i < j \leq m$.

(c) $Z_K = \sum_{i=0}^m Z_{B_i}$ (cf. also [29, (3.7)]).

(d) $A_iC'_t = A_iZ_K$ for any $A_i \subset |C'_t|$. In other words, C'_t is the canonical cycle of $|C'_t| = B_t$ (i.e. of $(M/B_t, p_t)$).

(e) For any $A_i \subset A$ the inequality $A_i \cdot C_t \leq 0$ holds.

Proof. a, b, c, d follows from the above construction. The proof of (e) is as follows. If $A_i \subset B_t$ then $A_iZ_{B_j} \leq 0$ for any $j \leq t$ by the definition of the numerical cycle, hence $A_iC_t \leq 0$. If $A_i \not\subset B_t$ then $A_iC_t = A_i(Z_K - C'_{t+1})$. Now, $A_iZ_K \leq 0$ (by the minimality of the resolution) and $A_iC'_{t+1} \geq 0$ (because $|C'_{t+1}| \subset B_t$). $\square$

2.12. The next result shows that the properties (d) and (e) characterize the elliptic sequence.

2.13. **Lemma.** *Assume that (X, p) is a numerical Gorenstein elliptic singularity.*

(a) *If a cycle $Z \geq 0$ satisfies $A_i(Z - Z_K) \geq 0$ for all $A_i \subset |Z|$, then in fact $A_i(Z - Z_K) = 0$ for all $A_i \subset |Z|$, and $Z \in \{C'_0, C'_1, \dots, C'_m, 0\}$.*

(b) *If a cycle $0 \leq Z \leq Z_K$ satisfies $A_i Z \leq 0$ for all $A_i \subset A$, then $Z \in \{0, C_0, C_1, \dots, C_m\}$.*

Proof. If $A_i(Z - Z_K) \geq 0$, then $Z(Z - Z_K) \geq 0$, hence $\chi(Z) \leq 0$. This together with (2.5) guarantees that $\chi(Z) = 0$ and $A_i(Z - Z_K) = 0$ for all $A_i \subset |Z|$. Now, if $|Z| = B_0$, then by the non-degeneracy of the intersection form $Z = Z_K$. If $|Z| \neq B_0$, then for $A_i \subset |Z|$ obviously $A_i(Z_K - Z) = 0$; for $A_i \not\subset |Z|$ one has $A_i(Z_K - Z) \leq 0$ (because $A_i Z_K \leq 0$ and $A_i Z \geq 0$). Hence $A_i(Z_K - Z) \leq 0$ for any A_i , therefore $Z_K - Z \geq Z_{B_0}$, hence $Z \leq Z_K - Z_{B_0}$. In particular, $|Z| \subset B_1$. If $|Z| = B_1$, then Z is the canonical cycle C'_1 of B_1 , otherwise, by the same argument as above $Z \leq Z_K - Z_{B_0} - Z_{B_1}$, hence $|Z| \subset B_2$. Continuing the process, (a) follows. For the second part, apply (a) for $Z_K - Z$. $\square$

Since $E \leq Z_{B_j} \leq Z_{num}$,

$$(2.14) \quad h^1(\mathcal{O}_{Z_{B_j}}) = 1 \quad \text{hence} \quad \chi(Z_{B_j}) = \chi(C_j) = \chi(C'_j) = 0 \quad \text{for all } 0 \leq j \leq m.$$

2.15. Since any Z_{B_i} is a numerical fundamental cycle, by [10] (proof of 4.1) or [12] (cf. also [29]), there is a “computation sequence” which can start with any of the irreducible exceptional divisors $A_j \subset B_i$ and ends with Z_{B_i} . This means that there exists a sequence of cycles $Z_0, \dots, Z_k$ with $Z_0 = A_j$, $Z_k = Z_{B_i}$, $Z_{l+1} = Z_l + A_{i_l}$ and $A_{i_l} Z_l > 0$ for $0 \leq l \leq k-1$. Moreover, for any two integers $0 \leq i < j \leq m$, since Z_{B_i} is a numerical cycle with $Z_{B_j} < Z_{B_i}$, there is a computation sequence which starts with Z_{B_j} and ends with Z_{B_i} . More precisely, there is a sequence $Z_0, \dots, Z_k$ with $Z_0 = Z_{B_j}$, $Z_k = Z_{B_i}$, $Z_{l+1} = Z_l + A_{i_l}$ and $A_{i_l} Z_l > 0$ for $0 \leq l \leq k-1$. Actually in this second case, since $h^1(\mathcal{O}_{Z_{B_i}}) = h^1(\mathcal{O}_{Z_{B_j}}) = 1$ (cf. 2.14), for any $0 \leq l \leq k-1$ the curve A_{i_l} is smooth rational and $A_{i_l} Z_l = 1$ (see also [12] (2.7)). We will call a sequence like this a “computation sequence which connects Z_{B_j} and Z_{B_i} ”.

Let D be a cycle with $DA_k = 0$ for any $A_k \subset B_i$, and consider a computation sequence $\{Z_l\}_l$ which connects Z_{B_j} and Z_{B_i} (where $j > i$). In the exact sequence $0 \rightarrow \mathcal{O}_{A_{i_l}}(-D - Z_l) \rightarrow \mathcal{O}_{D+Z_{l+1}} \rightarrow \mathcal{O}_{D+Z_l} \rightarrow 0$ ($0 \leq l \leq k-1$) the Chern number is $A_{i_l}(-D - Z_l) = -1$, hence the natural map

$$(2.16) \quad H^k(\mathcal{O}_{D+Z_{B_i}}) \rightarrow H^k(\mathcal{O}_{D+Z_{B_j}}) \quad (k = 1, 2) \text{ is an isomorphism.}$$

For example, if $D = C_{i-1}$ and $j = m$, then

$$(2.17) \quad H^k(\mathcal{O}_{C_i}) \rightarrow H^k(\mathcal{O}_{C_{i-1}+E}) \quad (1 \leq i \leq m; \quad k = 1, 2) \text{ is an isomorphism.}$$

Now consider the exact sequence

$$(2.18) \quad 0 \rightarrow \mathcal{O}_E(-C_{i-1}) \rightarrow \mathcal{O}_{C_{i-1}+E} \rightarrow \mathcal{O}_{C_{i-1}} \rightarrow 0.$$

Since E is 2-connected (see e.g. [17, (4.21)]), $\mathcal{O}_E(-C_{i-1})$ is trivial if and only if it has a non-zero section (see e.g. [17], page 82), and in this case $h^k(\mathcal{O}_E(-C_{i-1})) = 1$ for $k = 1, 2$. Therefore (2.17) and (2.18) imply $h^1(\mathcal{O}_{C_{i-1}}) \leq h^1(\mathcal{O}_{C_i}) \leq h^1(\mathcal{O}_{C_{i-1}}) + 1$ for $1 \leq i \leq m$. Since $h^1(\mathcal{O}_{C_0}) = 1$, one obtains S. S.-T. Yau's result ([29, (3.9)]):

$$(2.19) \quad h^1(\mathcal{O}_{C_i}) \leq i + 1 \text{ (for } 0 \leq i \leq m), \text{ hence } p_g = h^1(\mathcal{O}_{C_m}) \leq m + 1.$$

The next theorem gives some characterizations of the “extremal property” $p_g = m + 1$.

2.20. First characterization of $p_g = m + 1$. *Assume that (X, p) is a numerical Gorenstein elliptic singularity. Then the following facts are equivalent:*

- (a) $p_g = m + 1$;
- (b) $h^1(\mathcal{O}_{C_i}) = i + 1$ for all $0 \leq i \leq m$;
- (c) The line bundles $\mathcal{O}_E(-C_j)$ are trivial (in $\text{Pic}(E)$), and $H^0(\mathcal{O}_{C_j+E}) \rightarrow H^0(\mathcal{O}_{C_j})$ is surjective for any $0 \leq j \leq m - 1$;
- (d) $h^1(\mathcal{O}(-C_j)) = m - j$ for all $0 \leq j \leq m$;
- (e) $h^1(\mathcal{O}_{C'_j}) = m - j + 1$ for all $0 \leq j \leq m$;
- (f) For any $0 \leq j \leq m - 1$, there exists $f_j \in H^0(M, \mathcal{O}(-C_j))$, such that for any $A_l \subset B_{j+1}$ the vanishing order of f_j on A_l is exactly $m_{A_l}(C_j)$;
- (g) The line bundles $\mathcal{O}_{C'_{j+1}}(-C_j)$ are trivial for $0 \leq j \leq m - 1$;
- (h) The line bundles $\mathcal{O}_{C'_{j+1}}(-Z_{B_j})$ are trivial for $0 \leq j \leq m - 1$.

Proof. $a \Leftrightarrow b \Leftrightarrow c$ is just a reformulation of the above discussion (2.17-18-19). (Actually $a \Leftrightarrow c$ can be found in [25] (2.4).)

$d \Rightarrow a$ is easy: for $j = 0$ one has $m = h^1(\mathcal{O}(-Z_{num})) \stackrel{(2.3)}{=} h^1(\mathcal{O}) - h^1(\mathcal{O}_{Z_{num}}) \stackrel{(2.5)}{=} p_g - 1$. Conversely, for $(a, b, c) \Rightarrow d$, notice that $H^0(\mathcal{O}) \rightarrow H^0(\mathcal{O}_{C_m})$ is onto by (2.2), and $H^0(\mathcal{O}_{C_m}) \rightarrow H^0(\mathcal{O}_{C_j})$ is onto by part (c) and (2.17) applied several times. Therefore $H^0(\mathcal{O}) \rightarrow H^0(\mathcal{O}_{C_j})$ is onto, hence $h^1(\mathcal{O}(-C_j)) = h^1(\mathcal{O}) - h^1(\mathcal{O}_{C_j}) = m + 1 - j - 1$.

$e \Rightarrow a$ again is trivial (take $j = 0$). In order to prove $a \Rightarrow e$, we construct a sequence $\{Z_l\}_l$. We start with $Z_0 = Z_K$. If $Z_l > 0$ is already constructed, then we define $Z_{l+1} = Z_l - A_{i_l}$ as follows. If there is at least one $A_i \subset |Z_l|$ with $A_i(Z_l - Z_K) < 0$ then take for A_{i_l} one of these A_i 's. Otherwise A_{i_l} is an arbitrary $A_i \subset |Z_l|$. In this second case, by (2.13), $A_{i_l}(Z_l - Z_K) = 0$ and $Z_l \in \{C'_0, \dots, C'_m\}$. If $Z_l = 0$ then we stop.

Now, from the exact sequence $0 \rightarrow \mathcal{O}_{A_{i_l}}(-Z_{l+1}) \rightarrow \mathcal{O}_{Z_l} \rightarrow \mathcal{O}_{Z_{l+1}} \rightarrow 0$ and Serre duality one gets that $h^1(\mathcal{O}_{Z_l}) = h^1(\mathcal{O}_{Z_{l+1}})$ in the first case, and in second case $h^1(\mathcal{O}_{Z_{l+1}}) = h^1(\mathcal{O}_{Z_l}) - \epsilon_l$, where $\epsilon_l \in \{0, 1\}$. Since h^1 must drop exactly $m + 1$ times, we obtain that in the above sequence we reach all the cycles C'_j and at every time $\epsilon_l = 1$.

$a \Rightarrow f$ is proved in [29] (3.13). We present a short proof of it. Fix an arbitrary $A_k \subset B_{j+1}$. Consider a computation sequence $\{Z_l\}_l$ of $Z_{B_{j+1}}$ which starts with A_k (cf. 2.15). Since the Chern numbers $A_{i_l}(-Z_l - C_j) < 0$, using the sequences $0 \rightarrow \mathcal{O}_{A_{i_l}}(-Z_l - C_j) \rightarrow \mathcal{O}_{Z_{l+1}}(-C_j) \rightarrow \mathcal{O}_{Z_l}(-C_j) \rightarrow 0$, we obtain that $\beta : H^0(\mathcal{O}_{Z_{B_{j+1}}}(-C_j)) \rightarrow H^0(\mathcal{O}_{A_k}(-C_j))$ is injective. Now, $H^0(\mathcal{O}_{A_k}(-C_j)) = \mathbf{C}$. Indeed, if $A_k \approx \mathbf{P}^1$ then $H^0(\mathcal{O}_{A_k}(-C_j)) = H^0(\mathcal{O}_{\mathbf{P}^1})$. If $A_k \not\approx \mathbf{P}^1$ then $A_k = E$ (by 2.8), and $\mathcal{O}_{A_k}(-C_j)$ is trivial by (c). Now, considers the diagram:

$$\begin{array}{ccc}
H^0(\mathcal{O}(-C_j)) & \xrightarrow{\alpha_k} & H^0(\mathcal{O}_{A_k}(-C_j)) \\
\uparrow = & & \uparrow \beta \\
H^0(\mathcal{O}(-C_j)) & \xrightarrow{\alpha} & H^0(\mathcal{O}_{Z_{B_{j+1}}}(-C_j)) \rightarrow H^1(\mathcal{O}(-C_{j+1})) \rightarrow \\
& & H^1(\mathcal{O}(-C_j)) \rightarrow H^1(\mathcal{O}_{Z_{B_{j+1}}}(-C_j)) \rightarrow 0
\end{array}$$

By part (d): $h^1(\mathcal{O}(-C_t)) = m - t$, hence by dimension argument $h^1(\mathcal{O}_{Z_{B_{j+1}}}(-C_j)) \neq 0$. Since $\chi(\mathcal{O}_{Z_{B_{j+1}}}(-C_j)) = 0$ and β is injective, we obtain that, in fact, β is an isomorphism, and α is onto; hence α_k is onto as well. Finally, the surjectivity of α_k for all $A_k \subset B_{j+1}$ implies (f).

For $f \Rightarrow b$ notice that the image of the elements $1, f_0, \dots, f_{i-1}$ are linearly independent in $H^0(\mathcal{O}_{C_i})$, this together with (2.19) proves (b).

$f \Leftrightarrow g$. For any $A_l \subset B_{j+1}$ consider the composed map

$$H^0(M, \mathcal{O}(-C_j)) \xrightarrow{u} H^0(\mathcal{O}_{C'_{j+1}}(-C_j)) \xrightarrow{v} H^0(\mathcal{O}_{A_l}(-C_j)).$$

By (2.2) u is onto. Hence $v \circ u$ is onto if and only if v is onto. Notice that $h^0(\mathcal{O}_{A_l}(-C_j)) = 1$ always (if (f) is true then see the proof of $a \Rightarrow f$; if (g) is true, then $\mathcal{O}_{A_l}(-C_j)$ is a trivial line bundle).

$g \Leftrightarrow h$ is easy: $\mathcal{O}_{C'_{j+1}}(-C_j) = \otimes_{i \leq j} \mathcal{O}_{C'_{i+1}}(-Z_{B_i})|_{C'_{j+1}}$, and $\mathcal{O}_{C'_{j+1}}(-Z_{B_j}) = \mathcal{O}_{C'_{j+1}}(-C_j) \otimes \mathcal{O}_{C'_j}(-C_{j-1})|_{C'_{j+1}}^{-1}$. $\square$

2.21. Corollary. *Assume that (X, p) is an elliptic Gorenstein singularity.*

a) *Then $p_g = m + 1$ if and only if the line bundles $\mathcal{O}_{C'_{j+1}}(-Z_{B_j})$ are trivial for all $1 \leq j \leq m - 1$.*

b) *If $p_g = m + 1$, then $h^1(M, \mathcal{O}(-lZ_{num})) = h^1(Z_K, \mathcal{O}_{Z_K}(-lZ_{num})) = p_g - 1$ for any $l \geq 1$.*

Proof of (a). We have to prove that in the Gorenstein case the triviality of $\mathcal{O}_{C'_1}(-Z_{B_0})$ follows from the triviality of the other line bundles. Indeed, if all the other line bundles are trivial, then (2.20.a $\Leftarrow$ h), applied for $(M/B_1, p_1)$, gives $h^1(\mathcal{O}_{C'_1}) = m$. But the Gorenstein property implies that for any positive cycle $D < Z_K$ the strict inequality $h^1(\mathcal{O}_D) < h^1(\mathcal{O}_{Z_K}) = p_g$ holds (see e.g. [17], page 109). Hence $h^1(\mathcal{O}_{Z_K}) > h^1(\mathcal{O}_{C'_1}) = m$. Therefore $p_g = m + 1$. Now, we prove (b). Using (2.2) $h^1(\mathcal{O}(-lZ_{num})) = h^1(\mathcal{O}_{Z_K}(-lZ_{num})) = h^1(\mathcal{O}_{C'_1}(-lZ_{num}))$ for $l \geq 1$. But the line bundle $\mathcal{O}_{C'_1}(-lZ_{num})$ is trivial by (2.20.g), hence its first Betti-number is $p_g - 1$ by (2.20.e). $\square$


In the next sections, (2.21.b) will replace some vanishing theorems which were used in the classical case of minimally elliptic singularities (i.e. when $p_g - 1 = 0$).

2.22. Definition. *In the sequel, we call the line bundles $\mathcal{O}_{C'_{j+1}}(-Z_{B_j})$ ($0 \leq j \leq m - 1$) “obstruction line bundles”.*

The next example shows that the property $p_g = m + 1$ is not always true (even if we deal with Gorenstein singularities).

2.23. Example. Consider $(X_1, 0) = \{x^2 + y^3 + z^{18} = 0\} \subset (\mathbf{C}^3, 0)$ and $(X_2, 0) = \{z^2 =$

$y(x^4 + y^6)\} \subset (\mathbf{C}^3, 0)$ (cf. also [14, 9] and [29], page 291). Using the method of [9], Chapter II, (or the Appendix of [16]) one can verify that the minimal resolution graph in both cases is:



On the other hand, the geometric genus in the first case is $p_g = 3$, and in the second case is $p_g = 2$ (this can be verified using e.g. [13] or [16]).

Denote by A_0, A_1, A_2 the irreducible exceptional divisors (starting from left). Then $A_0 = E = Z_{B_2}$, $A_0 + A_1 = Z_{B_1}$, $A_0 + A_1 + A_2 = Z_{num}$, and $Z_K = 3A_0 + 2A_1 + A_2$ (hence $m = 2$). Moreover $\chi(Z_{num}) = 0$, hence the singularities are elliptic. It is very instructive to check property (2.20.f) for these examples. For a function $f : (X, 0) \rightarrow (\mathbf{C}, 0)$ we denote its divisor supported by A by $(f)_A := \sum_i m_{A_i}(f \circ \pi)A_i$.

In the first case, $Z_{num} = (z)_A$. This shows that in (2.20.f) (a possible choice is) $f_0 = z$ and $f_1 = z^2$. In the second case there is no function f_0 with $m_{A_0}(f_0) = m_{A_0}(Z_{num}) = 1$, hence (2.20.f) fails (in particular, there is no function f with $(f)_A = Z_{num}$).

3. Characterization of the trivial line bundles in $Pic(C'_j)$.

As usual, for any positive cycle D , we denote the isomorphism classes of invertible sheaves over $\mathcal{O}_D$ by $Pic(D)$ ($= H^1(\mathcal{O}_D^*)$). The kernel of the degree map $deg : Pic(D) \rightarrow \mathbf{Z}^{\#\{A_i | A_i \subset D\}}$ defined by $\mathcal{L} \rightarrow \{deg_{A_i} \mathcal{L}\}_i$ is denoted by $Pic^0(D)$.

If $D = E$ is a minimally elliptic cycle, then

$$(3.1) \quad \begin{cases} \mathcal{L} \in Pic^0(E) \text{ is trivial} \Leftrightarrow H^0(E, \mathcal{L}) \neq 0; \\ \text{moreover, if } \mathcal{L} \text{ is trivial then } h^0(E, \mathcal{L}) = 1. \end{cases}$$

This basically follows from the 2-connectivity of E (see e.g. [17] page 82). But, in general, it is not really easy to provide similar result for $Pic^0(D)$. Fortunately, in the case of an elliptic singularity, for any $0 \leq j \leq m$, $Pic^0(Z_{B_j})$ is as simple as $Pic^0(E)$. Indeed, consider the “exponential cohomology sequences” (see, e.g. [5] page 49) of the cycles Z_{B_j} and E . Then the natural maps $H^1(|Z_{B_j}|, \mathbf{Z}) \rightarrow H^1(|E|, \mathbf{Z})$ and $H^1(\mathcal{O}_{Z_{B_j}}) \rightarrow H^1(\mathcal{O}_E)$ are isomorphism (the first from 2.9, or by a similar argument; for the second use 2.16 for $D = 0$). Hence $Pic^0(Z_{B_j}) \rightarrow Pic^0(E)$ is an isomorphism as well. Moreover, if we consider a computation sequence $\{Z_l\}_l$ which connects $E = Z_{B_m}$ and Z_{B_j} (cf. 2.15), then from the exact sequences $0 \rightarrow \mathcal{O}_{A_{i_l}}(-Z_l) \otimes \mathcal{L} \rightarrow \mathcal{L}|_{Z_{l+1}} \rightarrow \mathcal{L}|_{Z_l} \rightarrow 0$ follows that $H^0(Z_{B_j}, \mathcal{L}) = H^0(E, \mathcal{L}|_E)$. Therefore, for any line bundle $\mathcal{L} \in Pic^0(Z_{B_j})$:

$$(3.2) \quad \begin{cases} h^0(Z_{B_j}, \mathcal{L}) \leq 1, \text{ and} \\ h^0(Z_{B_j}, \mathcal{L}) = 1 \text{ if and only if } \mathcal{L} \text{ is trivial.} \end{cases}$$

3.3. Theorem. *Assume that the elliptic numerical Gorenstein singularity (X, p) satisfies $p_g = m + 1$; and fix an integer $0 \leq j \leq m$. Then the followings are equivalent.*

- a) $\mathcal{L} \in \text{Pic}^0(C'_j)$ is trivial;
- b) $h^0(C'_j, \mathcal{L}) \geq m - j + 1$.

Actually, if $\mathcal{L}$ is trivial, then $h^0(C'_j, \mathcal{L})$ is exactly $m - j + 1$.

Proof. $a \Rightarrow b$ and the equality follow from (2.20; $a \Rightarrow e$) and (2.14). We prove $b \Rightarrow a$ by descending induction over j . If $j = m$, then $C'_j = E$, hence (a) follows from (3.1). Assume that $b \Rightarrow a$ is true for $j + 1$, and $h^0(C'_j, \mathcal{L}) \geq m - j + 1$. Consider the exact sequence

$$0 \rightarrow \mathcal{O}_{C'_{j+1}}(-Z_{B_j}) \otimes \mathcal{L} \rightarrow \mathcal{L} \rightarrow \mathcal{L}|_{Z_{B_j}} \rightarrow 0.$$

Then $h^0(\mathcal{L}|_{Z_{B_j}}) \leq 1$ by (3.2), hence $h^0 := h^0(\mathcal{O}_{C'_{j+1}}(-Z_{B_j}) \otimes \mathcal{L}) \geq m - j$. Hence, by the inductive step $\mathcal{O}_{C'_{j+1}}(-Z_{B_j}) \otimes \mathcal{L}$ is trivial in $\text{Pic}^0(C'_{j+1})$, and $h^0 = m - j$. Therefore $h^0(\mathcal{L}|_{Z_{B_j}})$ must be 1, $\mathcal{L}|_{Z_{B_j}}$ must be trivial, and $r : H^0(\mathcal{L}) \rightarrow H^0(\mathcal{L}|_{Z_{B_j}})$ onto. In particular, there exists $s \in H^0(\mathcal{L})$ such that $r(s)$ generates $\mathcal{O}(\mathcal{L}|_{Z_{B_j}})$. Since $|Z_{B_j}| = |C'_j|$, it follows that s generates the sections of $\mathcal{L}$ as well. $\square$

3.4. Corollary. *Assume that (X, p) is an elliptic Gorenstein singularity, and we fix an integer $0 \leq j \leq m$. Assume that $h^1(\mathcal{O}_{C'_j}) = m - j + 1$. Then $\mathcal{O}_{C'_j}(-C_{j-1})$ is trivial in $\text{Pic}^0(C'_j)$.*

For example, for $m = j$ we obtain that $\mathcal{O}_E(-C_{m-1})$ is trivial in $\text{Pic}^0(E)$ (the assumption $h^1(\mathcal{O}_E) = 1$ is automatically satisfied). For $j = m - 1$, if contracting B_{m-1} one has $p_g((M/B_{m-1}, p_{m-1})) = 2$, then $\mathcal{O}_{C'_{m-1}}(-C_{m-2})$ is trivial in $\text{Pic}^0(C'_{m-1})$.

Proof. First notice that contracting B_j we obtain a numerical Gorenstein singularity with canonical cycle C'_j (cf. 2.12), and with the “extremal property” $p_g(M/B_j, p_j) = \text{length of the elliptic sequence of } B_j = m - j + 1$. Therefore, (3.3) for this singularity gives: $\mathcal{L} \in \text{Pic}^0(C'_j)$ is trivial $\Leftrightarrow h^0(C'_j, \mathcal{L}) = m - j + 1$. But by Serre duality $h^0(\mathcal{O}_{C'_j}(-C_{j-1})) = h^1(\mathcal{O}_{C'_j}) = m - j + 1$. $\square$

3.5. Second characterization of $p_g = m + 1$. *Assume that (X, p) is a numerical Gorenstein elliptic singularity. Consider its minimal resolution $M \rightarrow X$ and the singularity $(M/B_j, p_j)$ obtained by the contraction of B_j ($0 \leq j \leq m$). Then (X, p) satisfies the extremal property $p_g = m + 1$ if and only if the singularities $(M/B_j, p_j)$ are Gorenstein for all $0 \leq j \leq m - 1$. (Notice that $(M/B_m, p_m)$ is automatically Gorenstein by 2.7.)*

Proof. “ $\Rightarrow$ ” First we prove that a numerical Gorenstein singularity (X, p) with $p_g = m + 1$ is Gorenstein. (This fact was proved by S. S.-T. Yau in [29, (3.11)]; we present here a very short proof in the spirit of [17, (4.21)]; cf. also with the papers of L. Bădescu [3, 4].)

Consider the line bundle $\mathcal{L} := \mathcal{O}_{Z_K}(Z_K + K) \in \text{Pic}^0(Z_K)$. By Serre duality $h^0(\mathcal{L}) = h^1(\mathcal{O}_{Z_K})$, hence $h^0(\mathcal{L}) = p_g = m + 1$. Therefore, by (3.3), $\mathcal{L}$ is trivial. Moreover, by (2.2) $h^1(\mathcal{O}(K)) = 0$, hence $H^0(M, \mathcal{O}(Z_K + K)) \rightarrow H^0(\mathcal{L}) \approx H^0(\mathcal{O}_{Z_K})$ is onto. Therefore, there is a global section of $\mathcal{O}(Z_K + K)$ which has no zeros in the neighbourhood of A , hence $Z_K + K$ is linearly equivalent to the zero cycle. But this is one of the characterizations of the Gorenstein property.

Now, M/B_j is an elliptic numerical Gorenstein singularity with numerical cycle C'_j and the length of its elliptic sequence $= h^1(\mathcal{O}_{C'_j}) = m - j + 1$ (cf. 2.11 and 2.20.e). Therefore, the above fact applied for the singularities M/B_j ends the proof.

“ $\Leftarrow$ ” By (2.20.h), it is enough to prove that the obstruction line bundles $\mathcal{L}_j := \mathcal{O}_{C'_j}(-Z_{B_{j-1}})$ are trivial for $1 \leq j \leq m$. We will prove this by induction. Assume that for a fixed $1 \leq j_0 \leq m$ the line bundles $\mathcal{L}_j$ are trivial for $j > j_0$ (if $j_0 = m$ then this assumption is vacuous). Since the singularity M/B_{j_0-1} is Gorenstein, by (2.20.h) and (2.21.a) (applied for M/B_{j_0-1}), the triviality of $\{\mathcal{L}_j\}_{j>j_0}$ implies the triviality of $\mathcal{L}_{j_0}$. $\square$

3.6. Examples. Assume that (X, p) is a numerical Gorenstein elliptic singularity.

(a) If $m = 1$ then $p_g \leq 2$, and:

$$p_g = 2 \Leftrightarrow \mathcal{O}_E(-Z_{num}) \text{ is trivial} \Leftrightarrow (X, p) \text{ is Gorenstein.}$$

Examples with $p_g = 1$ exists (actually this is the generic case!). E.g. take the minimal resolution of any Gorenstein singularity with $p_g = 2$ and $m = 1$. Then deform its analytic structure. Generically we obtain a non-Gorenstein singularity with $p_g = 1$ (cf. [12], page 1279).

(b) Assume that $m = 2$. Then in the numerical Gorenstein case we have two obstruction line bundles. For the generic analytic structure $p_g = 1$ as above. If (X, p) is Gorenstein, then $p_g = h^1(\mathcal{O}_{Z_K}) > h^1(\mathcal{O}_E) = 1$, hence $p_g \geq 2$. From (2.21.a), $p_g = 3$ if and only if $\mathcal{O}_E(-Z_{B_1})$ is trivial. From (3.4) $\mathcal{O}_E(-Z_{num} - Z_{B_1})$ is trivial, hence **if (X, p) is Gorenstein, then:**

$$p_g = 3 \Leftrightarrow \mathcal{O}_E(-Z_{num}) \text{ is trivial} \Leftrightarrow (M/B_1, p_1) \text{ is Gorenstein.}$$

4. Torsion properties of the “obstruction line bundles”.

The main result of this section is the following:

4.1. Theorem. *Consider the minimal resolution of an elliptic Gorenstein singularity (X, p) with elliptic sequence $\{Z_{B_j}\}_{j=0}^m$ ($m \geq 1$), and fix an integer $1 \leq k \leq m$.*

Assume that for $k+1 \leq j \leq m$ the “obstruction line bundles” $\mathcal{O}_{C'_j}(-Z_{B_{j-1}}) \in \text{Pic}^0(\mathcal{O}_{C'_j})$ are trivial. (If $k = m$ then this assumption is vacuous.) Then $\mathcal{O}_{C'_k}(-Z_{B_{k-1}}) \in \text{Pic}^0(\mathcal{O}_{C'_k})$ has finite order.

Actually, there exist integers $1 \leq l_r \leq k - r$ ($0 \leq r \leq k - 1$) such that for any $0 \leq r \leq k - 1$ the line bundle $\mathcal{O}_{C'_k}(l_0 \cdots l_r Z_{B_r})$ is trivial. In particular, the order of $\mathcal{O}_{C'_k}(-Z_{B_{k-1}})$ in $\text{Pic}^0(\mathcal{O}_{C'_k})$ is not greater than $k!$.

Proof. First assume that $k = 1$. Then $\mathcal{O}_{C'_1}(-Z_{B_0})$ is trivial by (2.21.a) and (2.20.a $\Leftrightarrow$ h). So, in the sequel we will assume that $k \geq 2$.

Since $(M/B_k, p_k)$ is a numerical Gorenstein singularity with canonical class C'_k (cf. 2.11.d), the assumption together with (2.20) (applied for this singularity) provide that:

$$(4.2) \quad h^1(\mathcal{O}_{C'_k}) = m - k + 1.$$

4.3. Lemma. For any $0 \leq r \leq t \leq k-2$ one has $h^1(\mathcal{O}(-C_t - Z_{B_r})) \geq m - k + 1$.

Proof. Using (2.2), Serre duality, (2.14), and in the last step $C'_{t+2} \geq C'_k$, one has: $h^1(\mathcal{O}(-C_{t+1})) \stackrel{2.2}{=} h^1(\mathcal{O}_{C'_{t+2}}(-C_{t+1})) \stackrel{SD}{=} h^0(\mathcal{O}_{C'_{t+2}}) \stackrel{2.14}{=} h^1(\mathcal{O}_{C'_{t+2}}) \geq h^1(\mathcal{O}_{C'_k})$. Hence, by (4.2):

$$(4.4) \quad h^1(\mathcal{O}(-C_{t+1})) \geq m - k + 1.$$

Now, consider a computation sequence $\{Z_l\}_l$ which connects $Z_{B_{t+1}}$ with Z_{B_r} (cf. 2.15) (i.e. $Z_{l+1} = Z_l + A_{i_l}$ where A_{i_l} is smooth rational curve with $A_{i_l}Z_l = 1$). In the exact sequence $0 \rightarrow \mathcal{O}(-C_t - Z_{l+1}) \rightarrow \mathcal{O}(-C_t - Z_l) \rightarrow \mathcal{O}_{A_{i_l}}(-C_t - Z_l) \rightarrow 0$ the Chern number $A_{i_l}(-C_t - Z_l)$ is $-A_{i_l}C_t - 1 \geq -1$ (cf. 2.11.e). Hence $H^1(\mathcal{O}(-C_t - Z_{l+1})) \rightarrow H^1(\mathcal{O}(-C_t - Z_l))$ is onto for any l , in particular, $h^1(\mathcal{O}(-C_t - Z_{B_r})) \geq h^1(\mathcal{O}(-C_t - Z_{B_{t+1}}))$. The lemma follows from this and (4.4). $\square$

Now consider the exact sequence:

$$(4.5) \quad 0 \rightarrow \mathcal{O}(-Z_K - Z_{B_r}) \rightarrow \mathcal{O}(-C_t - Z_{B_r}) \rightarrow \mathcal{O}_{C'_{t+1}}(-C_t - Z_{B_r}) \rightarrow 0.$$

4.6. Lemma. $H^1(\mathcal{O}(-Z_K - Z_{B_r})) = 0$.

Proof. Consider a computation sequence $\{Z_l\}_l$ which connects Z_{B_r} and $Z_{B_0} = Z_{num}$ (cf. 2.15). Consider the exact sequences:

$$0 \rightarrow \mathcal{O}(-Z_K - Z_{l+1}) \rightarrow \mathcal{O}(-Z_K - Z_l) \rightarrow \mathcal{O}_{A_{i_l}}(-Z_K - Z_l) \rightarrow 0$$

and notice that $A_{i_l}(-Z_K - Z_l) = -A_{i_l}Z_K - 1 \geq -1$ (cf. 2.1). Hence $H^1(\mathcal{O}(-Z_K - Z_{num})) \rightarrow H^1(\mathcal{O}(-Z_K - Z_{B_r}))$ is onto. But $H^1(\mathcal{O}(-Z_K - Z_{num})) = 0$ (by 2.2). $\square$

Now, from (4.5) and duality:

$$H^1(\mathcal{O}(-C_t - Z_{B_r})) = H^1(\mathcal{O}_{C'_{t+1}}(-C_t - Z_{B_r})) = H^0(\mathcal{O}_{C'_{t+1}}(C'_{t+1} + C_t + Z_{B_r} - Z_K)).$$

Hence (4.3) reads as follows: for $0 \leq r \leq t \leq k-2$:

$$h^0(\mathcal{O}_{C'_{t+1}}(C'_{t+1} + C_t + Z_{B_r} - Z_K)) \geq m - k + 1.$$

Now, for any positive cycle $0 < D \leq C'_{t+1}$ write $o(D) := \mathcal{O}_D(D + C_t + Z_{B_r} - Z_K)$. For any irreducible exceptional divisor $A_i \subset |D|$, one has the exact sequence:

$$0 \rightarrow o(D - A_i) \rightarrow o(D) \rightarrow \mathcal{O}_{A_i}(D + C_t + Z_{B_r} - Z_K) \rightarrow 0.$$

The Chern number $A_i(D + C_t + Z_{B_r} - Z_K) = A_i(D - Z_K)$, for $A_iZ_{B_r} = A_iC_t = 0$. Therefore, if $A_i(D - Z_K) < 0$, then $H^0(o(D - A_i)) \rightarrow H^0(o(D))$ is an isomorphism. In the sequel, we will construct a decreasing sequence of cycles as follows. The starting cycle is $D = C'_{t+1}$. If D is already constructed, and there exists $A_i \subset |D|$ with $A_i(D - Z_K) < 0$, then the next term is $D - A_i$. During this step $h^0(o(D))$ will stay constant.

If at a moment we reach a cycle $D > 0$ with the property $A_i(D - Z_K) \geq 0$ for all $A_i \subset |D|$, then by (2.13) in fact $A_i(D - Z_K) = 0$ for all $A_i \subset |D|$ and D must be one of the cycles C'_s . If there is a $A_i \subset |D|$ such that

$$r_i : H^0(\mathcal{O}_D(D + C_t + Z_{B_r} - Z_K)) \rightarrow H^0(\mathcal{O}_{A_i}(D + C_t + Z_{B_r} - Z_K))$$

is the trivial map, then the term following D is $D - A_i$ for this A_i . Hence, in this case again, $h^0(o(D)) = h^0(o(D - A_i))$. If the above map r_i is nontrivial for all $A_i \subset |D|$, then the term following D is $D - A_i$, where A_i arbitrary with $A_i \subset |D|$. In this later case r_i has rank one, hence $h^0(o(D))$ will decrease by one. If $D = 0$ we stop.

Since finally $h^0(o(D))$ must decrease to zero, and $h^0(o(C'_{t+1})) \geq m - k + 1$, $h^0(o(D))$ must decrease at least $m - k + 1$ times. Therefore, the first time when we decrease h^0 must happen for a cycle $D = C'_s$ with $s \leq k$ (i.e. the length of the sequence $\{C'_s, \dots, C'_m\}$ must be at least $m - k + 1$). Obviously $C'_s \leq C'_{t+1}$ (because we deal with cycles $D \leq C'_{t+1}$), hence: $s \geq t + 1$.

This shows that there exists $t + 1 \leq s \leq k$ such that

$$H^0(\mathcal{O}_{C'_s}(C'_s + C_t + Z_{B_r} - Z_K)) \rightarrow H^0(\mathcal{O}_{A_i}(C'_s + C_t + Z_{B_r} - Z_K)) = \mathbf{C}$$

is onto for all $A_i \subset |C'_s|$. In other words, $\mathcal{O}_{C'_s}(C'_s + C_t + Z_{B_r} - Z_K)$ is a trivial line bundle. Taking its restriction to C'_k , we obtain:

$$(4.7) \quad \begin{cases} \text{for any pair } (r, t) \text{ with } 0 \leq r \leq t \leq k - 2, \\ \text{there exists } s \text{ with } t + 1 \leq s \leq k, \text{ such that} \\ \mathcal{O}_{C'_k}(C'_s + C_t + Z_{B_r} - Z_K) \text{ is a trivial line bundle in } \text{Pic}^0(C'_k). \end{cases}$$

4.8. Lemma. *For any $0 \leq r \leq k - 1$, there exists l_r with $1 \leq l_r \leq k - r$, such that*

$$\mathcal{O}_{C'_k}(l_0 \cdots l_r Z_{B_r}) \text{ is trivial in } \text{Pic}^0(C'_k).$$

Proof. First we prove for $r = 0$. If there is at least one s in (4.7) with $s = t + 1$, then $C'_s + C_t - Z_K = 0$, hence $\mathcal{O}_{C'_k}(Z_{B_0})$ is trivial. Now, assume that $s \geq t + 2$ for any t . This means that for any $0 \leq t \leq k - 2$, there exists $t + 2 \leq s(t) \leq k$ such that:

$$(*_t) \quad \mathcal{O}_{C'_k}(Z_{B_0}) \approx \mathcal{O}(Z_{B_{t+1}} + \cdots + Z_{B_{s(t)-1}}).$$

Now, we set $t_1 := 0$. If $s(t_1) = k$ we stop; otherwise consider $t_2 := s(t_1) - 1$. If $s(t_2) = k$ we stop, otherwise we continue until we obtain $s(t_u) = k$ for some $u \geq 1$. If we multiply the identities $(*_t)$ for $t_1, \dots, t_u$, we obtain:

$$\mathcal{O}_{C'_k}(uZ_{B_0}) \approx \mathcal{O}_{C'_k}(Z_{B_1} + \cdots + Z_{B_{k-1}}).$$

Here u satisfies $1 \leq u \leq k - 1$ ($u = k - 1$ exactly when $s(t) = t + 2$ for every t , i.e. when we have to consider all the isomorphisms $(*_t)$). But from (3.4) and (4.2):

$$(4.9) \quad \mathcal{O}_{C'_k}(Z_{B_0} + \cdots + Z_{B_{k-1}}) \text{ is trivial,}$$

hence with the notation $l_0 := u + 1$ one has $\mathcal{O}_{C'_k}(l_0 Z_{B_0})$ is trivial.

Now, take $r = 1$ and consider again (4.7). If $s = t + 1$ for at least one t , then $\mathcal{O}_{C'_k}(Z_{B_1})$ is trivial. If $s \geq t + 2$ for all $1 \leq t \leq k - 2$, by a similar argument as above one obtains that $\mathcal{O}_{C'_k}(uZ_{B_1}) \approx \mathcal{O}_{C'_k}(Z_{B_2} + \cdots + Z_{B_{k-1}})$ for some $1 \leq u \leq k - 2$ ($u = k - 2$ occurs

exactly when we have to consider all the identities corresponding to $1 \leq t \leq k-2$). This together with (4.9) gives that $\mathcal{O}_{C'_k}(Z_{B_0} + (u+1)Z_{B_1})$ is trivial. Now consider its l_0^{th} power and take $l_1 := u+1$.

Similar argument can be used for all $0 \leq r \leq k-2$. For $r = k-1$, just take (4.9) again and take its $l_0 \cdots l_{k-2}$ -th power. $\square$

4.10. Example. (cf. 3.6.b) Assume that (X, p) is elliptic Gorenstein, and $m = 2$. Recall that $\mathcal{O}_E(-Z_{B_1} - Z_{num}) \approx \mathcal{O}_E$. The above theorem (for $m = k = 2$) says that the order of $\mathcal{O}_E(-Z_{num})$ divides 2, hence:

$$\mathcal{O}_E(-Z_{num}) \text{ is trivial} \Leftrightarrow p_g = 3; \quad \mathcal{O}_E(-Z_{num}) \text{ has order 2} \Leftrightarrow p_g = 2.$$

One of the main results of the present paper is the following:

4.11. Theorem. *Assume that (X, p) is an elliptic Gorenstein singularity with $H^1(A, \mathbf{Z}) = 0$. Then $p_g = m + 1 =$ the length of the elliptic sequence in the minimal resolution of (X, p) . In particular p_g is a topological invariant.*

Proof. If $H^1(A, \mathbf{Z}) = 0$, then by the exponential exact sequence ([5], page 49) $Pic^0(D) \approx H^1(\mathcal{O}_D)$ for any positive cycle D , in particular, it is torsion free. Therefore, (4.1) (as an inductive step) proves that all the obstruction line bundles are trivial. Hence $p_g = m + 1$ by (2.20). $\square$

4.12. Remark. If $m \geq 1$, then in (4.11) both assumptions are necessary. Without Gorenstein condition p_g generically is one ([12, page 1279]). Moreover, Example (2.23) shows that $p_g < m + 1$ can occur if $H^1(A, \mathbf{Z}) \neq 0$.

4.13. Corollary. *For a Gorenstein singularity (X, p) with $H^1(A, \mathbf{Z}) = 0$ the following facts are equivalent:*

$$p_g = 2 \Leftrightarrow \chi(Z_{num}) = 0 \text{ and } Z_K = Z_{num} + E.$$

(Notice that the right hand side is completely topological.)

Proof. Indeed, if (X, p) is Gorenstein with $p_g = 2$, then $h^1(Z_{num})$ cannot be zero (because of 2.4), cannot be greater than or equal to two (that would imply that $Z_{num} = Z_K$, which characterise the minimally elliptic singularities, cf. 2.7), hence it is one. Therefore (X, p) is elliptic. The rest follows from the above results. $\square$

4.14. Remark. S. S.-T. Yau in [29], Theorem B, proved that a Gorenstein singularity with $p_g = 2$ is elliptic. Moreover, in [27], he proved also that a *hypersurface* singularity with $H^1(A, \mathbf{Z}) = 0$ and $p_g = 2$ satisfies $m = 1$. His proof (of this second fact) is based on the classification of all possible dual resolution graphs of hypersurface singularities with $p_g = 2$ (250 cases).

Notice that the “second characterization” (3.5) gives:

4.15. Corollary. *If (X, p) is an elliptic Gorenstein singularity with $H^1(A, \mathbf{Z}) = 0$ then the singularities $(M/B_j, p_j)$ are Gorenstein for all $0 \leq j \leq m$.*

5. The multiplicity.

5.1. Assume that (X, p) is an elliptic Gorenstein singularity with $p_g = m + 1$. In this section we will prove that its multiplicity $\text{mult}(X, p)$ is a topological invariant, in fact depends only on Z_{num}^2 .

First notice that by (2.20.f) there exists $f_0 \in H^0(M, \mathcal{O}(-Z_{\text{num}}))$ such that for any $A_l \subset B_1$, the order of vanishing of f_0 on A_l is exactly $m_{A_l}(Z_{\text{num}})$. In particular:

$$(5.2) \quad H^0(M, \mathcal{O}(-Z_{\text{num}})) \rightarrow H^0(\mathcal{O}_{A_l}(-Z_{\text{num}})) = \mathbf{C} \text{ is onto for all } A_l \subset B_1.$$

By (2.2), if $|\mathcal{O}_{Z_K}(-Z_{\text{num}})|$ is free, then $|\mathcal{O}(-Z_{\text{num}})|$ is free too. By [21, page 426], if $|\mathcal{O}(-Z_{\text{num}})|$ is free, then $m_p \mathcal{O}_M = \mathcal{O}(-Z_{\text{num}})$, in which case $\text{mult}(X, p) = -Z_{\text{num}}^2$. In particular:

$$(5.3) \quad Z_{\text{num}}^2 = -1 \Rightarrow -Z_{\text{num}}^2 \neq \text{mult}(X, p) \Rightarrow |\mathcal{O}_{Z_K}(-Z_{\text{num}})| \text{ is not free.}$$

If $|\mathcal{O}(-Z_{\text{num}})|$ is not free, then $\text{mult}(X, p) > -Z_{\text{num}}^2$. E.g. if $m_p \mathcal{O}_M = m_Q \mathcal{O}(-Z_{\text{num}})$, where m_Q is the maximal ideal of a smooth point Q of A , then $\text{mult}(X, p) = -Z_{\text{num}}^2 + 1$.

The next theorem generalizes Laufer's result about the multiplicity of minimally elliptic singularities [12].

5.4. Theorem. *Assume that (X, p) is an elliptic Gorenstein singularity with $p_g = m + 1$. (If $m = 0$ or $H^1(A, \mathbf{Z}) = 0$ then the last assumption is satisfied.) Then:*

If $Z_{\text{num}}^2 \leq -2$, then $m_p \mathcal{O}_M = \mathcal{O}(-Z_{\text{num}})$, hence $\text{mult}(X, p) = -Z_{\text{num}}^2$;

If $Z_{\text{num}}^2 = -1$, then $m_p \mathcal{O}_M = m_Q \mathcal{O}(-Z_{\text{num}})$ for some smooth point Q of A , and $\text{mult}(X, p) = 2$.

We prove theorem (5.4) in several steps. First we prove the converse of (5.3). For simplicity, in the sequel we will write $\mathcal{L} := \mathcal{O}_{Z_K}(-Z_{\text{num}})$.

5.5. Proposition. *Assume that $|\mathcal{L}|$ is not free. Then $Z_{\text{num}}^2 = -1$.*

Proof. By (5.2), all the basepoints must lie outside of B_1 . Since all the connected components of $A \setminus B_1$ are trees whose vertices correspond to smooth rational curves (cf. 2.8), we can fix a basepoint Q such that Q is "as close to B_1 as possible". More precisely, we will fix a basepoint Q such that there exist irreducible exceptional divisors $\{A'_i\}_{i=0}^l$ with

$$(5.6) \quad \begin{cases} A'_i \not\subset B_1 \text{ and } A'_i \text{ is smooth rational curve } (0 \leq i \leq l); \\ Q \in A'_0, A'_i A'_{i+1} = 1 \text{ for } 0 \leq i \leq l-1, \text{ and } A'_l B_1 = 1; \\ \text{there is no basepoint of } |\mathcal{L}| \text{ on } \cup_{i>0} A'_i. \end{cases}$$

With this choice of Q , $A'_0 Z_{\text{num}}$ cannot be zero. Indeed, since Q is a basepoint, the evaluation map $H^0(\mathcal{L}) \xrightarrow{\alpha} H^0(\mathcal{O}_{A'_0}(-Z_{\text{num}})) \xrightarrow{e} \mathbf{C}_Q$ is trivial. If, by assumption, $A'_0 Z_{\text{num}} = 0$ then e must be an isomorphism, hence α must be trivial as well. This shows that $A'_0 \cap A'_1$ is also a basepoint, which contradicts the above choice of Q in (5.6). For later reference:

$$(5.7) \quad A'_0 Z_{\text{num}} < 0.$$

Since the evaluation map $H^0(Z_K, \mathcal{L}) \rightarrow \mathbf{C}_Q$ is zero, from the exact sequence $0 \rightarrow m_Q \mathcal{L} \rightarrow \mathcal{L} \rightarrow \mathbf{C}_Q \rightarrow 0$ and (2.21.b), one gets $h^1(m_Q \mathcal{L}) = h^1(\mathcal{L}) + 1 = p_g$.

In the next paragraph, we follow the beginning of the proof of (4.23) of [17]. Take $0 < D \leq Z_K$ minimal with the property $h^1(D, m_Q \mathcal{L}|_D) = p_g$. Minimality means:

$$h^1(D - A_i, m_Q \mathcal{L}|_{D-A_i}) \leq p_g - 1 \text{ for any } A_i \subset |D|.$$

Let K_i be the kernel of $m_Q \mathcal{L}|_D \rightarrow m_Q \mathcal{L}|_{D-A_i}$. Outside Q , clearly $K_i = \mathcal{L} \otimes \mathcal{O}_{A_i}(-D + A_i)$. Actually, if $Q \in |D - A_i|$, then even at Q one has $K_i = \mathcal{L} \otimes \mathcal{O}_{A_i}(-D + A_i)$. If $Q \in |D|$ but $Q \notin |D - A_i|$ (i.e. if $Q \in A_i \setminus \{\text{other components of } D\}$ and $m_{A_i}(D) = 1$), then $K_i = \mathcal{L} \otimes m_Q \mathcal{O}_{A_i}(-D + A_i)$. We define $\delta_i := 0$ in the first two cases, and $\delta_i := 1$ in the third case.

Now, the minimality of D implies that for all i : $h^1(D, K_i) \neq 0$, in other words:

$$(5.8) \quad A_i \mathcal{L} + A_i(Z_K - D) \leq \delta_i \text{ for all } A_i \subset |D|.$$

5.9. Lemma. *The situation when $A_i \mathcal{L} + A_i(Z_K - D) = 0$ for all $A_i \subset |D|$ cannot occur.*

Proof. Assume the contrary. Since $A_i \mathcal{L} \geq 0$ for all A_i , and $\chi(D) \geq 0$, the relation

$$0 \leq D(Z_K - D) = \sum m_{A_i}(D) A_i(Z_K - D) = - \sum m_{A_i}(D) A_i \mathcal{L} \leq 0$$

says that $A_i \mathcal{L} = A_i(Z_K - D) = 0$ for all $A_i \subset |D|$. In particular, $D = C'_s$ for some $0 \leq s \leq m$ (by 2.13), and $D \mathcal{L} = C'_s(-Z_{num}) = 0$. Since $C'_0(-Z_{num}) = -Z_{num}^2 \neq 0$, we must have $s \geq 1$, i.e. $|D| \subset B_1$. On the other hand, (5.2) guarantees that $Q \notin B_1$, hence $Q \notin |D|$. Therefore, $m_Q \mathcal{L}|_D = \mathcal{O}_{C'_s}(-Z_{num})$ for some $s \geq 1$. Using (3.3) we obtain $h^1(\mathcal{O}_{C'_s}(-Z_{num})) = h^0(\mathcal{O}_{C'_s}(-Z_{num})) \leq m - s + 1 \leq p_g - 1$, which contradicts $h^1(D, m_Q \mathcal{L}|_D) = p_g$. $\square$

The above lemma implies that there exists $A_{i_0} \subset |D|$ with $A_{i_0} \mathcal{L} + A_{i_0}(Z_K - D) = \delta_{i_0} = 1$. (In particular, either $Q = A_{i_0} \cap A'_0$ and $A'_0 \not\subset |D|$, or $A_{i_0} = A'_0$, cf. 5.6. The first case will be eliminated later.) Summing (5.8) over all components of D (recall $m_{A_{i_0}}(D) = 1$) we obtain $D \mathcal{L} + D(Z_K - D) = 1$. Since $D \mathcal{L} \geq 0$ (by the definition of Z_{num}) and $D(Z_K - D)$ is an *even* non-negative integer (by Riemann-Roch), we get:

$$(5.10) \quad D \mathcal{L} = 1 \text{ and } D(Z_K - D) = 0.$$

If we write $D = \sum D_i$, where $\{|D_i|\}_i$ are the the connected components of $|D|$, since $\chi(D)=0$ and $\chi(D_i) \geq 0$ for all i (cf. 2.5), we must have $\chi(D_i) = 0$ for all i . This shows that no D_i lies in $A \setminus B_1$ (for all the components of $A \setminus B_1$ support rational singularities whose positive cycles have $\chi > 0$, cf. 2.8 and 2.4). Therefore, all the exceptional divisors $\{A'_i\}_{i=0}^l$ defined in (5.6) must be components of $|D|$ and $A'_0 = A_{i_0}$. Then (5.7) reads as

$$(5.11) \quad A_{i_0} Z_{num} < 0.$$

Now, $A_j \mathcal{L} \geq 0$ for all $A_j \subset |D|$ and $D \mathcal{L} = 1$ (cf. 5.10), hence there is exactly one index j_0 with $A_{j_0} Z_{num} \leq -1$. (5.11) guarantees that this j_0 is exactly i_0 . Therefore, we get:


$$\begin{aligned} A_{i_0} Z_{num} &= -1, & A_{i_0}(Z_K - D) &= 0; \\ A_j Z_{num} &= 0, & A_j(Z_K - D) &= 0 \quad \text{for } j \neq i_0; & A_j &\subset |D|. \end{aligned}$$

Again by (2.13): $D = C'_s$ for some $0 \leq s \leq m$. Since $Q \in |D|$ but $Q \notin B_1$, the only possibility is $D = Z_K$. This means that $1 = D\mathcal{L} = Z_K(-Z_{num}) = -Z_{num}^2$, which ends the proof of (5.5). $\square$

5.12. The special case $Z_{num}^2 = -1$. All the possible dual (minimal) resolution graphs of elliptic numerical Gorenstein singularities with $Z_{num}^2 = -1$ are classified. This is done by S. S.-T. Yau (in [26], (2.3) and Table 1) based on Laufer's classification of the minimal resolution graphs of the minimally elliptic singularities with $E^2 = -1$ ([12], Table 1.)

5.13. Proposition. [26, 12] *The minimal resolution graph of a numerical Gorenstein elliptic singularity with $Z_{num}^2 = -1$ has the form $A = B_1 \cup A_0$, where A_0 is a smooth rational curve with self-intersection -2 , and it is unique with $A_0 Z_{num} = -1$. Moreover, $m_{A_0}(Z_{num}) = m_{A_0}(Z_K) = 1$. B_1 supports a singularity with $Z_{B_1}^2 = -1$. The curve A_0 intersects only one irreducible exceptional divisor $\tilde{A}$ of B_1 , and it intersects this exactly in one point. The curve $\tilde{A}$ is the unique component of B_1 with the property $\tilde{A} Z_{B_1} = -1$.*

Therefore, by induction, $A = |E| \cup A_{m-1} \cup \dots \cup A_0$, where all curves $\{A_i\}_{i=0}^{m-1}$ are smooth rational curves with self-intersection -2 , $E^2 = -1$, $Z_{B_j} = E + \sum_{i \geq j} A_i$, and the dual graph has the following form:



Now, Theorem 5.4 follows from the above discussions and the next proposition.

5.14. Proposition. *Assume that (X, p) is numerical Gorenstein elliptic singularity with $p_g = m + 1$ ($m > 0$) and $Z_{num}^2 = -1$. Consider its minimal resolution graph and write $A = B_1 \cup A_0$ (cf. 5.13). Then $|\mathcal{O}(-Z_{num})|$ has a unique basepoint $Q \in A_0 \setminus B_1$, and $m_p \mathcal{O}_M = m_Q \mathcal{O}_M(-Z_{num})$.*

The *proof* is similar to [12] (3.13), and it is left to the reader.

6. The Hilbert–Samuel function and the embedding dimension.

Assume that (X, p) is an elliptic Gorenstein singularity with $p_g = m + 1$. Consider its minimal resolution and set $\mathcal{L} := \mathcal{O}_{Z_K}(-Z_{num})$ and $d := -Z_{num}^2$.

6.1. Theorem. *If $d \geq 3$ then $\oplus_{k \geq 0} H^0(Z_K, \mathcal{L}^{\otimes k})$ is generated by elements of degree $k = 1$.*

Proof. We will follow – and modify – the proof of the corresponding statement for the minimally elliptic singularities (as it is presented in [17], page 114; i.e. “Castelnuovo’s free pencil trick”).

By (5.4) there exists $s_0 \in H^0(\mathcal{L})$ with $m_{A_i}(s_0) = m_{A_i}(Z_{num})$ for all A_i . Its divisor has form $Z_{num} + D$, where D is an effective Cartier divisor with $|D| \cap A$ finite, and $A_i D = A_i \mathcal{L}$. Obviously $\mathcal{O}_{Z_K}(D) \rightarrow \mathcal{L}$ (given by $f \mapsto f s_0$) is an isomorphism. We use another generic section $s \in H^0(\mathcal{L})$ to identify $\mathcal{L}$ with $\mathcal{O}_{Z_K}$ near D , hence we will write $\mathcal{O}_D$ instead of $\mathcal{L}|_D = \mathcal{O}_D(D)$. By construction, $\mathcal{O}_D$ is an Artinian scheme of degree d . Consider the

exact sequence $0 \rightarrow \mathcal{O}_{Z_K} \rightarrow \mathcal{L} \rightarrow \mathcal{O}_D \rightarrow 0$, where the second arrow is the multiplication by s_0 , (and the third is the “division by s near D ”). Its long cohomology exact sequence is:

$$0 \rightarrow H^0(\mathcal{O}_{Z_K}) \xrightarrow{s_0} H^0(\mathcal{L}) \xrightarrow{r_1} \mathcal{O}_D \rightarrow H^1(\mathcal{O}_{Z_K}) \rightarrow H^1(\mathcal{L}) \rightarrow 0.$$

Since $h^1(\mathcal{O}_{Z_K}) = p_g$ and $h^1(\mathcal{L}) = p_g - 1$ (cf. 2.21.b), the rank of r_1 is $d - 1$.

In the following lemmas $D' \subset D$ is an arbitrary subscheme of length $d - 1$. Let $\mathcal{I}_{D'} \subset \mathcal{O}_{Z_K}$ be its ideal sheaf. If $\mathcal{I}_D \subset \mathcal{O}_{Z_K}$ is the ideal sheaf of D , then $\mathcal{I}_D \subset \mathcal{I}_{D'}$ and $\mathcal{I}_{D'}/\mathcal{I}_D = \mathbf{C}_Q$ for some point $Q \in A$.

6.2. Lemma. $h^1(\mathcal{I}_{D'}\mathcal{L}) = p_g - 1$.

Proof. The inclusion $\mathcal{I}_{D'}\mathcal{O}_{Z_K}(D) \subset \mathcal{O}_{Z_K}(D) = \mathcal{L}$ guarantee that $\mathcal{I}_{D'}\mathcal{L}$ is torsion free. Hence the exact sequence $0 \rightarrow \mathcal{O}_{Z_K} \xrightarrow{s_0} \mathcal{I}_{D'}\mathcal{L} \rightarrow \mathbf{C}_Q \rightarrow 0$ does not split. Hence in the exact sequence $0 \rightarrow \text{Hom}(\mathcal{I}_{D'}\mathcal{L}, \mathcal{O}_{Z_K}) \rightarrow \text{Hom}(\mathcal{O}_{Z_K}, \mathcal{O}_{Z_K}) \xrightarrow{\alpha} \text{Ext}^1(\mathbf{C}_Q, \mathcal{O}_{Z_K}) = \mathbf{C}$ the map α is not trivial. Hence by Serre duality $h^1(\mathcal{I}_{D'}\mathcal{L}) = h^1(\mathcal{O}_{Z_K}) - 1$. $\square$

6.3. Lemma. $H^0(\mathcal{L}) \rightarrow \mathcal{O}_{D'}$ is surjective for any subscheme $D' \subset D$ of length $d - 1$.

Proof. Use the cohomology exact sequence of $0 \rightarrow \mathcal{I}_{D'}\mathcal{L} \rightarrow \mathcal{L} \rightarrow \mathcal{O}_{D'} \rightarrow 0$ and $h^1(\mathcal{I}_{D'}\mathcal{L}) = h^1(\mathcal{L})$ (by 6.2. and 2.21.b). $\square$

Now, by similar arguments as in [17], pages 114-115, Theorem 6.1 and Theorem D (of the introduction) follows. $\square$

6.4. Remarks.

(a) **The cases** $d = -Z_{num}^2 = 1, 2$. Similarly to the minimally elliptic singularities (cf. e.g. [17], page 116), one can write a generator set of the $\mathcal{O}_X$ -algebra $R := \bigoplus_{k \geq 0} H^0(\mathcal{O}(-kZ_{num}))$ in the case $d \leq 2$ as well. If $d = 2$, then R is generated by m_p in degree 1 and an element $y \in m_p \setminus m_p^2$ in degree two. If $d = 1$, then R is generated by m_p in degree 1, by an element $y \in m_p \setminus m_p^2$ in degree 2, and $z \in m_p \setminus (m_p^2, y)$ in degree 3.

(b) Moreover (cf. [12]), if $Z_{num}^2 = -4$, then (X, p) is a (tangential) complete intersection; if $Z_{num}^2 \leq -5$, then (X, p) is not a complete intersection.

(c) Using similar arguments as in this section, one can verify that for any normal surface singularity $\text{emb dim}(X, p) \geq \text{mult}(X, p) + \min_{D > 0} \chi(D)$. Actually, for any elliptic Gorenstein singularity $\text{emb dim}(X, p) = \text{mult}(X, p)$ provided that $\text{mult}(X, p) \geq 3$, and (obviously) $\text{emb dim}(X, p) = 3$ if $\text{mult}(X, p) = 2$. In particular, for elliptic hypersurface singularities $\text{mult}(X, p) \leq 3$.

(d) S. S.-T. Yau computed the multiplicity and the Hilbert–Samuel function for many different situations. His basic assumption was the much stronger $Z_E^2 \leq -2$ for the first case and $Z_E^2 \leq -3$ for the second case. (For details see his series of papers listed in the References.)

(e) If $Z_{num}^2 = -1$, then (5.13) shows that the dual resolution graph is a “Kodaira graph” (cf. e.g. [6], 2.7). Moreover, by (5.14), the maximal ideal cycle is exactly Z_{num} . Hence, by [6] (2.9.1), (X, p) is a Kodaira singularity.

References

- [1] M. Artin: Some numerical criteria for contractibility of curves on algebraic surfaces. Amer. J. of Math., **84**, 485-496 (1962)
- [2] M. Artin: On isolated rational singularities of surfaces. Amer. J. of Math., **88**, 129-136 (1966)
- [3] L. Bădescu: Applications of the Grothendieck duality theory to the study of normal isolated singularities. Rev. Roum. Math. Pures et Appl., Tome **XXIV**, No 5, 673-689 (1979)
- [4] L. Bădescu: Dualising divisors of two-dimensional singularities. Rev. Roum. Math. Pures et Appl., Tome **XXV**, No 5, 695-707 (1980)
- [5] W. Barth, C. Peters and A. Van de Ven: Compact complex surfaces. Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band 4, A Series of Modern Surveys in Mathematics, Springer-Verlag 1984.
- [6] U. Karras: On Pencils of Curves and Deformations of Minimally Elliptic Singularities. Math. Ann., **247**, 43-65 (1980)
- [7] U. Karras: Methoden zur Berechnung algebraischer Invarianten und zur Konstruktion von Deformationen normaler Flächensingularitäten. Habilitationsschrift, Dortmund 1981
- [8] V. S. Kulikov: Degenerate elliptic curves and deformation of uni- and bimodal singularities. Func. Anal. Appl., **9**, 69-70 (1975)
- [9] H. Laufer: Normal two-dimensional singularities. Annals of Math. Studies, **71**, Princeton University Press, 1971
- [10] H. Laufer: On rational singularities. Amer. J. of Math., **94**, 597-608 (1972)
- [11] H. Laufer: Taut two-dimensional singularities. Math. Ann., **205**, 131-164 (1973)
- [12] H. Laufer: On minimally elliptic singularities. Amer. J. of Math., **99**, 1257-1295 (1977)
- [13] H. Laufer: On μ for surface singularities. Proceedings of Symposia in Pure Math., **30**, 45-49 (1977)
- [14] H. Laufer: The multiplicity of isolated two-dimensional hypersurface singularities. Transactions of the AMS, **302** Number 2, 489-496 (1987)
- [15] W. D. Neumann: A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves. Transactions of the AMS, **268** Number 2, 299-344 (1981)
- [16] A. Némethi: The signature of $f(x, y) + z^N$. To appear in Proceedings of Real and Complex Singularities (C.T.C. Wall's 60th birthday meeting), Liverpool (England), August 1996.
- [17] M. Reid: Chapters on Algebraic Surfaces. In: Complex Algebraic Geometry, IAS/Park City Mathematical Series, Volume **3** (J. Kollár editor) 1997.
- [18] J.-P. Serre: Groupes Algébriques et Corps de Classes. Hermann, Paris 1959.

- [19] J. Stevens: Elliptic Surface Singularities and Smoothings of Curves. *Math. Ann.*, **267**, 239-247 (1984)
- [20] J. Stevens: Kulikov singularities. Thesis, Leiden 1985
- [21] Ph. Wagreich: Elliptic singularities of surfaces. *Amer. J. of Math.*, **92**, 419-454 (1970)
- [22] J. M. Wahl: Vanishing Theorems for Resolutions of Surface Singularities. *Inventiones math.*, **31**, 17-41 (1975)
- [23] S. S.-T. Yau: On almost minimally elliptic singularities. *Bulletin of the AMS*, **83** Number 3, 362-364 (1977)
- [24] S. S.-T. Yau: Normal singularities of surfaces. *Proc. of Symposia in Pure Math.*, **32**, 195-198 (1978)
- [25] S. S.-T. Yau: Normal two-dimensional elliptic singularities. *Transactions of AMS*, **254**, 117-134 (1979)
- [26] S. S.-T. Yau: Hypersurface dual graphs of normal singularities of surfaces. *Amer. J. of Math.*, **101**, 781-812 (1979)
- [27] S. S.-T. Yau: Gorenstein singularities with geometric genus equal to one. *Amer. J. of Math.*, **101**, 813-854 (1979)
- [28] S. S.-T. Yau: On strongly elliptic singularities. *Amer. J. of Math.*, **101**, 855-884 (1979)
- [29] S. S.-T. Yau: On maximally elliptic singularities. *Transactions of the AMS*, **257** Number 2, 269-329 (1980)